


This listing of claims will replace all prior versions, and listings, of claims in the application.

**Listing of Claims:**

1. (Previously presented) A compound of Formula (I):



Formula (I)

wherein

X, Y and Z are independently selected from the group consisting of CH and N; wherein m is an integer from 2 to 5; wherein X, Y and Z include at least one CH atom and at least one N atom; and, wherein a N atom may simultaneously occupy only the X and Z positions;

R<sub>1</sub> is selected from the group consisting of hydrogen and NH<sub>2</sub>; and,

R<sub>2</sub> is selected from the group consisting of phenyl, wherein phenyl is substituted with one substituent selected from the group consisting of halogen and heterocyclyl, and 1,4-benzodioxinyl; or a pharmaceutically acceptable salt thereof.

2. (Original) The compound of claim 1 wherein X, Y and Z are independently selected from the group consisting of CH and N; wherein m is 3; wherein X, Y and Z include at least one CH atom and at least one N atom; wherein a N atom may simultaneously occupy only the X and Z positions; wherein the heteroaryl ring thus formed is selected from the group consisting of pyridinyl and pyrazinyl; wherein pyridinyl is attached to the triazine ring at the 3 or 4 position of the pyridine ring; and, wherein pyrazinyl is attached to the triazine ring at the 6 position of the pyrazine ring.
3. (Original) The compound of claim 1 wherein R<sub>2</sub> is selected from the group consisting of phenyl (wherein phenyl is substituted with one substituent selected from the group consisting of chlorine and 4-morpholinyl) and 1,4-benzodioxinyl.


4. (Original) The compound of claim 1 wherein the compound of Formula (I) is selected from a compound wherein m is 3; and, wherein X, Y, Z, R<sub>1</sub> and R<sub>2</sub> are dependently selected from:

X	Y	Z	R <sub>1</sub>	R <sub>2</sub>
N	CH	CH	H	3-Cl-Ph;
CH	N	CH	H	3-Cl-Ph;
N	CH	N	H	3-Cl-Ph;
CH	N	CH	NH <sub>2</sub>	3-Cl-Ph;
N	CH	CH	H	2,3-dihydro-1,4-benzodioxin-6-yl; or,
N	CH	CH	H	4-(4-morpholinyl)Ph.

5. (Original) A composition comprising a compound of claim 1 and a pharmaceutically appropriate carrier.
6. (Canceled)
7. (Original) A method for preparing a composition comprising mixing a compound of claim 1 and a pharmaceutically appropriate carrier.
8. (Canceled)
9. (Canceled)
10. (Canceled)
11. (Canceled)
12. (Canceled)
13. (Canceled)
14. (Canceled)
15. (Canceled)
16. (Canceled)
17. (Canceled)

18. (Canceled)
19. (Canceled)
20. (Canceled)
21. (Canceled)
22. (Canceled)
23. (Canceled)
24. (Canceled)
25. (Canceled)
26. (Canceled)
27. (Canceled)
28. (Canceled)
29. (Canceled)
30. (Canceled)
31. (Canceled)
32. (Canceled)
33. (Canceled)
34. (Canceled)
35. (Canceled)
36. (Canceled)

37. (Previously presented) A method for treating human melanoma in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of Formula (I):



Formula (I)

wherein


X, Y and Z are independently selected from the group consisting of CH and N; wherein m is an integer from 2 to 5; wherein X, Y and Z include at least one CH atom and at least one N atom; and, wherein a N atom may simultaneously occupy only the X and Z positions;

R<sub>1</sub> is selected from the group consisting of hydrogen and NH<sub>2</sub>; and,

R<sub>2</sub> is selected from the group consisting of phenyl, (wherein phenyl is substituted with one substituent selected from the group consisting of halogen and heterocyclyl), and 1,4-benzodioxinyl;

or a pharmaceutically acceptable salt thereof.

38. (Previously presented) The method of claim 37 wherein the therapeutically effective amount of the compound of Claim 1 is from about 0.001 mg/kg/day to about 300 mg/kg/day.
39. (Previously presented) A method for treating rheumatoid arthritis in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of Formula (I):



Formula (I)

wherein

X, Y and Z are independently selected from the group consisting of CH and N; wherein m is an integer from 2 to 5; wherein X, Y and Z include at least one CH atom and at least one N atom; and, wherein a N atom may simultaneously occupy only the X and Z positions;

R1 is selected from the group consisting of hydrogen and NH<sub>2</sub>; and,

R2 is selected from the group consisting of phenyl, (wherein phenyl is substituted with one substituent selected from the group consisting of halogen and heterocyclyl), and 1,4-benzodioxinyl;

or a pharmaceutically acceptable salt thereof.

40. (Previously presented) The method of claim 39 wherein the therapeutically effective amount of the compound of Claim 1 is from about 0.001 mg/kg/day to about 300 mg/kg/day.